

Reference for Solicitation Entitled:

STRATEGIC IMPROVEMENTS TO RAPID SIMPLE POINT-OF-CARE TESTS (POCT) FOR SEXUALLY TRANSMITTED DISEASES (STDS)

1. [Peeling RW](#), et al “Rapid tests for sexually transmitted infections (STIs): the way forward. “ *Sex Transm Infect.* 2006. 82 Suppl 5:v1-6
2. Jackman, J et al. ”Minding the Gap: An approach to determine critical drivers in the development of Point of Care diagnostics.” *Point Care.* 2012. 11(2):130-139.
3. [Hsieh Y-H.](#) et al. “Perceptions on Point-of-Care Tests for Sexually Transmitted Infections - Comparison between Frontline Clinicians and Professionals in Industry. *Point Care.* 2012 .11(2):126-129.
4. Rompalo, A.M. et al. “Point-of-care tests for sexually transmissible infections: what do 'end users' want?” *Sexual health.* 2013.10(6): 541-5.
5. [Braverman P. K.](#)”Sexually transmitted diseases in adolescents.” *Med Clin North Am.* 2000 84(4):869-89, vi-vii
6. Hsieh Yu H. et al “What qualities are most important to making a point of care test desirable for clinicians and others offering sexually transmitted infection testing?” *PLoS One.* 2011. 6(4):e19263.
7. Peeling RW. “Diagnostics in a digital age: an opportunity to strengthen health systems and improve health outcomes.” *Int. Health.* 2015. 7(6):384-9.
8. [Vickerman P.](#) et al. Sensitivity requirements for the point of care diagnosis of Chlamydia trachomatis and Neisseria gonorrhoeae in women. *Sex Transm Infect.* 2003. 79(5):363-7.
9. www.cdc.gov/std/chlamydia/stdfact-chlamydia-detailed.htm (accessed 15 Jan 2019).